

Wyoming Basin & Water Supply Outlook Report April 1, 2022






Photo courtesy of Wyoming Snow Survey

Basin Outlook Reports

And

Federal - State - Private Cooperative Snow Surveys

For more Wyoming water supply information, contact:

Jim Fahey - Hydrologist 100 East "B" Street, Casper, WY 82601 (307) 233-6787 james.fahey@usda.gov

How forecasts are made

Most of the annual streamflow in the western United States originates as snowfall that has accumulated in the mountains during the winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Measurements of snow water equivalent at selected manual snow courses and automated SNOTEL sites, along with precipitation, antecedent streamflow, and indices of the El Niño / Southern Oscillation are used in computerized statistical and simulation models to prepare runoff forecasts. Unless otherwise specified, all forecasts are for flows that would occur naturally without any upstream influences.

Forecasts of any kind, of course, are not perfect. Streamflow forecast uncertainty arises from three primary sources: (1) uncertain knowledge of future weather conditions, (2) uncertainty in the forecasting procedure, and (3) errors in the data. The forecast, therefore, must be interpreted not as a single value but rather as a range of values with specific probabilities of occurrence. The middle of the range is expressed by the 50% exceedance probability forecast, for which there is a 50% chance that the actual flow will be above, and a 50% chance that the actual flow will be below, this value. To describe the expected range around this 50% value, four other forecasts are provided, two smaller values (90% and 70% exceedance probability) and two larger values (30%, and 10% exceedance probability). For example, there is a 90% chance that the actual flow will be more than the 90% exceedance probability forecast. The others can be interpreted similarly.

The wider the spread among these values, the more uncertain the forecast. As the season progresses, forecasts become more accurate, primarily because a greater portion of the future weather conditions become known; this is reflected by a narrowing of the range around the 50% exceedance probability forecast. Users should take this uncertainty into consideration when making operational decisions by selecting forecasts corresponding to the level of risk they are willing to assume about the amount of water to be expected. If users anticipate receiving a lesser supply of water, or if they wish to increase their chances of having an adequate supply of water for their operations, they may want to base their decisions on the 90% or 70% exceedance probability forecasts, or something in between. On the other hand, if users are concerned about receiving too much water (for example, threat of flooding), they may want to base their decisions on the 30% or 10% exceedance probability forecasts, or something in between. Regardless of the forecast value users choose for operations, they should be prepared to deal with either more or less water. (Users should remember that even if the 90% exceedance probability forecast is used, there is still a 10% chance of receiving less than this amount.) By using the exceedance probability information, users can easily determine the chances of receiving more or less water.

<u>Note</u>: The <u>median</u> is the official normal for snowpack (SWE), precipitation, reservoir storage, and streamflow calculations. Please refer to the **Appendix** of this report for more detailed information.


The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers. If you believe you experienced discrimination when obtaining services from USDA, participating in a USDA program, or participating in a program that receives financial assistance from USDA, you may file a complaint with USDA. Information about how to file a discrimination complaint is available from the Office of the Assistant Secretary for Civil Rights. USDA prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex (including gender identity and expression), marital status, familial status, parental status, religion, sexual orientation, political beliefs, genetic information, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibite bases apply to all programs.) To file a complaint of discrimination, complete, sign, and mail a program discrimination complaint form, available at any USDA office location or online at www.ascr.usda.gov, or write to: USDA Office of the Assistant Secretary for Civil Rights 1400 Independence Avenue, SW. Washington, DC 20250-9410 Or call toll free at (866) 632-9992 (voice) to obtain additional information, the appropriate office or to request documents. Individuals who are deaf, hard of hearing, or have speech disabilities may contact USDA through the Federal Relay service at (800) 877-8339 or (800) 845-6136 (in Spanish). USDA is an equal opportunity provider, employer, and lender. Persons with disabilities who require alternative means for communication of program information (e.g., Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

Summary

- Wyoming snowpack and/or snow water equivalents (SWEs)were below median by April 1st.
- Precipitation totals across Wyoming for March were **below** median. Water year precipitation totals lowered to **below** median.
- Overall reservoir storages for late March continue to be below median.
- Stream flow snowmelt volumes during April through July across Wyoming are forecasted to be generally **below** median.

Snowpack/SWEs


Snow water equivalents (SWEs) across Wyoming for April 1st were around 75% of median. SWEs in the Laramie and Tongue Watersheds were the highest at near 90% of median, while SWEs in the Belle Fourche and Cheyenne River Basins were the lowest at near 25% of median. Last year, SWEs across the state were 95% of median. (For complete *tabular data*, see attachment to report)




Map 1. Wyoming SWEs—April 1, 2022.

Precipitation

Basin precipitation across Wyoming was near 75% of median during March. The Lower North Platte Watershed had the <u>highest</u> precipitation totals for the month at near 150% of median. The Madison Headwaters, Belle Fourche, and Cheyenne River Basins had the <u>lowest</u> precipitation amounts at 40 to 50% of median. Water year precipitation (October - March) is currently about 90% of median. (For complete *tabular data*, see attachment to report)



Map 2. Current monthly precipitation by basin.



Map 3. Water year to date precipitation by basin.





Chart 1. Departure from median precipitation (water year).

Reservoirs

Reservoirs across Wyoming were averaging near 64% of capacity—down from 74% of capacity last year. Overall reservoir storages for late March are below median at 87% (101% last year). The Upper Green Basin had reservoir storages at 111% of median, while the Snake River Basin had reservoir storages at 30% of median. (For complete **tabular data**, see attachment to report)



Map 4. Reservoir storage (teacup diagram) by basin.

Stream Flows

April-July snowmelt runoff stream flow volumes across the state are expected to be **below** median at around 75%. The <u>highest</u> forecasted stream flows due to snowmelt are across the Wind and Laramie Basins at around 90% of median. The <u>lowest</u> snowmelt runoff volumes are expected across the Snake, Upper Green, and Lower Green Drainages between 60% and 70% of median. (For complete **tabular data**, see attachment to report)





Chart 2. Percent of Median (50% exceedance) for major forecast points.


April 1, 2022





Snake River Basin


- The overall Snake River Basin SWE is near 70% of median.
- Last month's precipitation for the Snake River Basin was near **50**% of median. Water-year-to-date precipitation is near **85**% of median.
- \bullet Current reservoir storage is near 30% of median for the two main reservoirs in the basin.
- The streamflow forecasts for April through July are **below** median (70%) for this basin. Jackson Lake inflows are also forecasted to be 70% of median.













Madison Headwaters Basin

- The overall basin SWE is around 70% of median.
- Last month's precipitation for the basin was near 40% of median. Water-year-to-date precipitation is around 80% of median.





No reservoir data for the basin.


There are no streamflow forecast points for the basin.




Yellowstone River Basin

- The overall Yellowstone River Basin SWE is near 75% of median.
- Last month's precipitation for the Yellowstone River Basin was near 60% of median. Water-year-to-date precipitation is near 85% of median.
- The 50% exceedance forecasts for April through July are **below** median (75%) for this basin. Clarks Fork near Belfry is forecasted to have flows at 78% of median.

Yellowstone River Basin **Snow Water Equivalent** 35.0 30.0 25.0 20.0 15.0 10.0 5.0 0.0 Feb Jan Mar Apr --x-- Current - Median Maximum Minimum




No reservoir data for the basin.








Wind River Basin

- The overall Wind River Basin SWE is near 85% of median.
- Last month's precipitation for the Wind River Basin was near 75% of median. Water-year-to-date precipitation is around 90% of median.
- Current reservoir storage remains close to 105% of median for three main reservoirs in the basin.
- The streamflow forecasts for April through July are below median (88%) for this basin. Boysen Reservoir inflows are expected to be 77% of median.















Bighorn River Basin

- The overall Bighorn River Basin SWE is near 80% of median.
- Last month's precipitation for the Bighorn River Basin was near 75% of median. Water-year-to-date precipitation is also about 90% of median.
- Current reservoir storage is near 100% of median for one main reservoir in the basin.
- The 50% exceedance forecasts for April through July are below median (82%) for this basin.















Shoshone River Basin

- The overall Shoshone River Basin SWE is around 75% of median.
- Last month's precipitation for the Shoshone River Basin was close to 60% of median. Water-year-to-date precipitation is around 85% of median.
- Current reservoir storage is near 90% of median for one main reservoir in the basin.
- Streamflow forecasts for April through July are **below** median (77%) for this basin. Buffalo Bill Reservoir inflows are expected to be 76% of median













Powder River Basin

- The overall Powder River Basin SWE is near 80% of median.
- Last month's precipitation for the Powder River Basin was near **75**% of median. Water-year-to-date precipitation is near **95**% of median.
- The 50% exceedance forecasts for April through July continue to be below median (68%) for this basin.

Snow Water Equivalent 15.0 10.0 5.0 Jan Feb Mar Apr Current — Median Maximum — Minimum




No reservoir data for the basin.








Tongue River Basin

- The overall Tongue River Basin SWE is near 90% of median.
- Last month's precipitation for the Tongue River Basin was at 60% of median. Water-year-to-date precipitation is close to 100% of median.
- Current reservoir storage remains near 105% of median for one main reservoir in the basin.
- The 50% exceedance forecasts for April through July continue to be **below** median (**80**%) for this basin. Tongue River Reservoir is forecasted to have inflows at **74**% of median.














Belle Fourche River Basin


- The overall Belle Fourche River Basin SWE is close to 25% of median.
- Last month's precipitation for the Belle Fourche River Basin was near 50% of median. Water-year-to-date precipitation is around 80% of median.
- Current reservoir storage is near 85% of median for three main reservoirs in the basin.









There are no streamflow forecast points for the basin.





Cheyenne River Basin

- The overall Cheyenne River Basin SWE is near 25% of median.
- Last month's precipitation for the Cheyenne River Basin was near 45% of median. Water-year-to-date precipitation is around 70% of median.
- ullet Current reservoir storage remains near 90% of median for three main reservoirs in the basin.
- The 50% exceedance forecasts for April through July are well below median (42%) for this basin. Pactola Reservoir inflows are forecasted to be only 30% of median.















Upper North Platte River Basin

- The overall Upper North Platte River Basin SWE is around 85% of median.
- Last month's precipitation for the Upper North Platte River Basin was near 90% of median. Water-year-to-date precipitation is around 95% of median.
- Current reservoir storage remains near 80% of median for two main reservoirs in the basin.
- Streamflow forecasts for April through July have dropped to **below** median (87%) for this basin. Seminoe Reservoir inflows are forecasted to be 85% of median.















Lower North Platte River Basin

- The overall Lower North Platte River Basin SWE is near 90% of median.
- Last month's precipitation for the Lower North Platte River Basin was near 150% of median. Water-year-to-date precipitation remains around 115% of median.
- Current reservoir storage remains near 95% of median for three main reservoirs in the basin.
- The 50% exceedance forecasts for April through July remain **below** median (86%) for this basin. La Prele Creek near Douglas is forecasted to have flows at 95% of median.















Laramie River Basin


- The overall Laramie River Basin SWE is around 90% of median.
- Last month's precipitation for the Laramie River Basin was near 100% of median. Water-year-to-date precipitation is close to 100% of median.
- ullet Current reservoir storage remains around 80% of median for one main reservoir in the basin.
- Streamflow forecasts for April through July have dropped **below** median (91%) for this basin. Laramie River near Woods Landing is expected to have flows at 100% of median.













Sweetwater River Basin

- The overall Sweetwater River Basin SWE is close to 70% of median.
- Last month's precipitation for the Sweetwater River Basin was close to 65% of median. Water-year-to-date precipitation is at 80% of median.
- Streamflow forecasts for April through July remain below median (73%) for this basin.

Sweetwater River Basin Snow Water Equivalent 30.0 25.0 20.0 2 15.0 20.0 10.0 5.0 0.0 Apr Feb Jan Mar - Current — 👉 – Median Maximum Minimum

No reservoir data for the basin.








South Platte River Basin (WY)

- The overall South Platte River Basin SWE is around 90% of median.
- Last month's precipitation for the South Platte River Basin was near 115% of median. Water-year-to-date precipitation remains near 95% of median.

South Platte River Basin **Snow Water Equivalent** 12.0 10.0 8.0 6.0 4.0 2.0 0.0 Feb Jan Mar Apr - \leftrightarrow - Current Median Maximum Minimum




No reservoir data for the basin.


There are $\underline{\text{no}}$ streamflow forecast points for the basin.





Little Snake River Basin

- The overall Little Snake River Basin SWE is near 85% of median.
- Last month's precipitation for the Little Snake River Basin was near 90% of median. Water-year-to-date precipitation remains close to 95% of median.
- Current reservoir storage is near 55% of median for one main reservoir in the basin.
- The 50% exceedance forecasts for April through July remain below median (80%) for this basin.















Upper Green River Basin

- The overall Upper Green River Basin SWE is near 70% of median.
- Last month's precipitation for the Upper Green River Basin was near **60**% of median. Water-year-to-date precipitation is around **85**% of median.
- Current reservoir storage is remains close to 110% of median for two main reservoirs in the basin.
- Streamflow forecasts for April through July have dropped to **well below** median (66%) for this basin. Fontenelle Reservoir inflows are expected to be 60% of median.















Lower Green River Basin


- The overall Lower Green River Basin SWE is near 80% of median.
- Last month's precipitation for the Lower Green River Basin was near **80**% of median. Water-year-to-date precipitation remains around **100**% of median.
- Current reservoir storage remains 90 to 95% of median for four main reservoirs in the basin.
- Streamflow forecasts for April through July have dropped to **well below** median (61%) for this basin. Flaming Gorge Reservoir inflows are forecasted to be 51% of median.















Upper Bear River Basin

- The overall Upper Bear River Basin SWE is close to 75% of median.
- Last month's precipitation for the Upper Bear River Basin was near **70**% of median. Water-year-to-date precipitation is around **95**% of median.
- ullet Current reservoir storage is near 45% of median for two main reservoirs in the basin.
- The 50% exceedance forecasts for April through July have dropped to **well below** median (68%) for this basin. Bear River above Woodruff Narrows Reservoir is expected to have flows at 68% of median.









Appendix

MEDIAN INFORMATION

Transitioning from 1981 – 2010 Averages to 1991 – 2020 Medians

Starting January 2022, the NRCS will use the 30-year **median** as the official normal for snowpack (SWE), precipitation, reservoir storage, and streamflow calculations. The National Water and Climate Center (NWCC) will continue to publish and distribute 30-year <u>averages</u> for alternate normal calculations.

The 30-yr reference period for median and normal calculations has also been recently updated from 1981-2010 to 1991-2020.

Please refer to this NWCC website or more information about the significant changes in data and forecast computations:

https://www.nrcs.usda.gov/wps/portal/wcc/home/snowClimateMonitoring/30YearNormals/

Topics include:

- 1991 2020 Median/Averages Overview
- Calculation Methods
- Differences Between 1991-2020 and Previous Normals
- Median vs. Average
- Retrieving 1991-2020 Normals

For specific seasonal <u>streamflow</u> normal comparisons for NRCS forecasted stations, please refer to:

https://www.wcc.nrcs.usda.gov/ftpref/support/srvo_norms_comps/

LINKS (for more information/graphics)

National Water Climate Center (NWCC)

➤ Interactive maps featuring current conditions of snow, precipitation, reservoir storages:

https://www.nrcs.usda.gov/wps/portal/wcc/home/quicklinks/predefinedMaps/

Water Resources Data System and State Climate Office (WRDS)

Clearinghouse of hydrological and climatological data for the State of Wyoming:
http://www.wrds.uwyo.edu/

USGS WaterWatch

> Tools and products to monitor streamflow, runoff, drought, and floods:

https://waterwatch.usgs.gov/index.php

Wyoming Basin Outlook Report National Resources Conservation Service Casper, Wyoming

Issued by:

Released by:

Terry Cosby (Chief)
U.S.D.A.
Natural Resources Conservation Service
Washington D.C.

Astrid Martinez State Conservationist N R C S Casper, Wyoming

The Following Agencies and Organizations Cooperate with the Natural Resources Conservation Service with Snow Surveys and/or with Data:

FEDERAL:

United States Department of the Interior (National Park Service)

United States Department of the Interior (Bureau of Reclamation)

United States Department of Agriculture (Forest Service)

United States Department of Commerce NOAA (National Weather Service)

STATE:

The Wyoming State Engineer's Office

The University of Wyoming

LOCAL:

The City of Cheyenne